

Volatile constituents of essential oils isolated from leaf and rhizome of *Zingiber cassumunar* Roxb.

Md. Nazrul Islam Bhuiyan, Jasim Uddin Chowdhury and Jaripa Begum

BCSIR Laboratories, Chittagong Cantonment, Chittagong 4220, Bangladesh.

Article Info

Received: 14 May 2008

Accepted: 23 May 2008

Available online: 23 May 2008

Keywords:

Essential oil

GC-MS

Sabinene

Triquinacene, 1,4-bis (methoxy)

Zingiber cassumunar

Number of Table: 1

Number of Refs: 21

Correspondence: MNIB

e-mail: nazrul119@yahoo.com

Abstract

The composition of the essential oil of *Zingiber cassumunar* Roxb. from Bangladesh was examined by gas chromatography mass spectroscopy (GC-MS). Sixty-four components were identified in leaf oil and 32 components were identified in the rhizome oil, accounting for 94.60% and 98.56% of the total yields, respectively. The main component in leaf oil was sabinene (14.99%), β -pinene (14.32%), caryophyllene oxide (13.85%) and caryophyllene (9.47%). On the other hand, rhizome oil contained triquinacene 1,4-bis (methoxy) (26.47%), (Z)-ocimene (21.97%) and terpinen-4-ol (18.45%). The compositions of both oils varied qualitatively and quantitatively.

Introduction

Zingiber cassumunar Roxb. synonyms, *Zingiber purpureum* Rosc., *Zingiber cliffordiae*, *Zingiber montanum* (J. König) Theilade (Family: Zingiberaceae) is used in folk medicine for the treatment of conditions such as inflammation, sprains, rheumatism, muscular pain, wounds and asthma, and as a mosquito repellent, a carminative, a mild laxative and an antidysenteric agent, cough and used as a cleansing solution for skin diseases (Oliveros, 1996). *Z. cassumunar* essential oil was found to exhibit absolute fungitoxic activity (Tripathi et al., 2008). A number of pure compounds isolated from the plants have been shown to possess antimicrobial (Wasuwat et al., 1989; Giwanon et al., 2000; Habsah et al., 2000),

topical and oral anti-inflammatory (Masuda and Jitoe, 1994; Kuroyanagi et al., 1980; Tuntiwachwuttikul et al., 1980, 1981; Panthong et al., 1990; Ozaki et al., 1991; Pongprayoon et al., 1996; Panthong et al., 1997; Jeenapongsa et al., 2003), antioxidative activity (Masuda and Jitoe, 1994; Kuroyanagi et al., 1980; Tuntiwachwuttikul et al., 1980, 1981; Kanjanapothi et al., 1987; Ozaki et al., 1991; Pongprayoon et al., 1997; Habsah et al., 2000) and antihistaminic effect (Piomrat et al., 1986) as well as activity as a smooth-muscle relaxant (Kanjanapothi et al., 1987). The rhizome oil of *Z. cassumunar* Roxb. from Malaysia was found to exhibit high activity against yeasts (Bin et al., 2003). Rhizomes of *Z. purpureum* Rosc. yield about 1.84 % of a colorless volatile oil which contains terpinen-4-ol (45.44 %) as its main

constituent (Oliveros, 1996). The essential oil obtained by hydrodistillation contained about 25 % of these phenylbutanoids whereas the oil obtained by extraction with light petroleum had about 46 %, with sabinene and terpinen-4-ol, *trans*-1-(3,4-dimethoxyphenyl)but-1-ene, *trans*-1-(3,4-dimethoxyphenyl)butadiene and *trans*-4-(3,4-dimethoxyphenyl)but-3-ene-1-yl acetate as the main constituents (Taroeno et al., 1991). The main active chemical constituents of the rhizome oil are sabinene (27-34%), γ -terpinene (6-8%), α -terpinene (4-5%), terpinen-4-ol (30-35%), and (E)-1-(3,4-dimethoxyphenyl)butadiene (DMPBD) (12-19%) (Pongprayoon et al., 1997). Jitoe et al. (1994) and Masuda and Jitoe, (1994) reported novel antioxidants, cassumunarin A, B, and C and Kishore and Dwivedi (1992) also reported a potential fungitoxic agent like Zerumbone from *Zingiber Cassumunar*. Regarding *Z. cassumunar*, there is no work available in our country. The present study deals with the investigation of the chemical components in leaf and rhizome oils of *Z. cassumunar* grown in Bangladesh.

Material and Methods

Plant material: The plant materials of *Zingiber cassumunar* were collected from the plants grown in the campus of BCSIR Laboratory, Chittagong during June 2007. One-voucher specimen (Y-694) was deposited in the herbarium of BCSIR Laboratory, Chittagong.

Extraction of essential oil: Samples of leaf was harvested from healthy, well-grown, two-year-old plants. Freshly leaf harvested samples (700 g) and the fresh rhizomes (600 g or 73 g of dry rhizomes) was sliced into small pieces and ground in a blender. The material was subjected to hydrodistillation using a modified Clevenger-type glass apparatus for 4 hour for isolation of oils separately from the two parts. The oil samples were stored at 0°C in air-tight containers after drying them over anhydrous sodium sulfate, filtered and concentrated under reduced pressure at room temperature to obtain the essential oil for GC-MS analyses.

GC-MS analysis: The essential oil from leaf and rhizome of *Zingiber cassumunar* was analyzed by GC-MS electron impact ionization (EI) method on GC-17A gas chromatograph (Shimadzu) coupled

to a GC-MS QP 5050A mass spectrometer (Shimadzu); fused silica capillary column (30 m x 2.5 mm; 0.25 μ m film thickness), coated with DB-1 (J&W); column temperature 100°C (2 min) to 250°C at the rate of 3°C/min; carrier gas, helium at constant pressure of 90 Kpa. Acquisition parameters full scan; scan range 40-350 amu.

Identification of the compounds: Compound identification was done by comparing the NIST library data of the peaks with those reported in literature, mass spectra of the peaks with literature data. Percentage composition was computed from GC peak areas on BP-I column without applying correction factors.

Results and Discussion

Essential oil from the leaf and rhizome of *Z. cassumunar* from Bangladesh were analyzed by GC-MS. The oil yields were 0.60% and 0.95% respectively. Table I reported the composition of the leaf and rhizome oil of *Z. cassumunar* from Bangladesh. According to GC-MS analysis under the conditions described above, sabinene was detected as the main component (14.99%) of the leaf essential oil. The remaining constituents, including β -pinene (14.32%), caryophyllene oxide (13.85%), caryophyllene (9.47%), γ -pinene (6.31%), methyl p-methoxycinnamate (5.02%), triquinacene, 1,4, bis(methoxy) (3.79%) and camphene (3.56%), were present at low concentrations. The rhizome oil was rich in triquinacene, 1,4-bis (methoxy) (26.47%), (Z)-ocimene (21.97%), terpinen-4-ol (18.45%), γ -terpinene (3.86%), β -phellandrene (3.49%) and cis-sabinenehydrate (3.00%). A major part of the oil consists of monoterpenes with sabinene and terpinen-4-ol as main common constituents of all the reported oils in the world including ours (Taroeno et al., 1991; Oliveros, 1996; Pongprayoon et al., 1997). *Z. cassumunar* oil was found to contain sabinene and terpinen-4-ol as its major constituents, which are no doubt responsible for its antimicrobial activities (Giwanon et al., 2000; Wasuwat et al., 1989). The study reveals that composition of two oils differs from the earlier reports and may, therefore be treated as different chemotypes. On the basis of above fact it may be concluded that *Z. cassumunar*, growing widely in Bangladesh, may be utilized as a source for the isolation of natural sabinene and triquinacene, 1,4-

bis (methoxy) respectively. In this study, the essential oil of *Z. cassumunar* has been extracted

Table I. Constituents of essential oil from the leaf and rhizome of *Zingiber cassumunar*

	Name of constituents in leaves	%	Name of constituents in rhizome	%
1.	(-)-Spathulenol	0.15	α -Thujene	0.70
2.	(E)-Ocimene	1.21	α -Pinene	2.30
3.	1,6,10-Dodecatriene, 7,11-dimethyl-3-methylene, (Z)	1.24	Camphene	0.36
4.	3,4,5-Trimethoxybenzylchloride	0.11	(Z)-Ocimene	21.97
5.	3-Cyclobexen-1-one, 3(hydroxymethyl)-6-(1-methylethyl)	0.17	β -Pinene	2.55
6.	3-Cyclohexene-1-methanol	0.18	β -myrcene	1.58
7.	Terpinen-4-ol	0.33	4-terpinyl acetate	2.10
8.	5-Caranol, trans	0.12	m-Cymene	0.46
9.	5-Nonaol,-5-methyl	0.20	β -Phellandrene	3.49
10.	7-Hexadecenal	0.11	γ -Terpinene	3.86
11.	7-Oxabicyclo (2.2.1) hept-5-en-2-one	0.10	Cis-Sabinenehydrate	3.00
12.	Apiol	1.40	2-Carene	0.77
13.	Aromadendrene oxide	0.12	Borneol	0.24
14.	Asaraldehyde	0.63	Terpinen-4-ol	18.45
15.	β -Bisabolene	0.19	Terpinyl acetate	1.10
16.	β -Elemene	0.17	Trans-Piperitol	0.34
17.	Benzen-1-methyl, 4-(1-methylethyl)	0.58	Bornyl acetate	0.29
18.	Bergamotol, z- α -trans	0.08	1,6,10-Dodecatrien,7,11-dimethyl-3-methylene (Z)	0.32
19.	β -Linalool	0.39	Germacrene D	0.15
20.	β -Myrcene	1.46	γ -Selinene	0.27
21.	Borneol	2.96	α -Selinene	0.12
22.	Bornyl acetate	0.63	α -Bergamotene	0.41
23.	β -Phellandrene	1.04	β -Bisabolene	0.13
24.	β -Pinene	14.32	β -Sesquiphellandrene	2.45
25.	β -Sesquiphellandrene	1.21	Methyleugenol	2.07
26.	Camphene	3.56	Megastigmastriene	0.33
27.	Caryophyllene	9.47	Lachnophyllum ester	0.59
28.	Caryophyllene oxide	13.85	2-Allyl-1,4-dimethoxy-3-methyl benzene	1.74
29.	Cedrene	0.40	Triquinacene,1,4-bis (methoxy)	26.47
30.	Chamigrene	0.15	δ -Cadinene	0.13
31.	Cholestan-3-ol, 2-methylene-(3B,5L)	0.88	Juniper camphor	0.44
32.	cis-Bicyclo (4.4.0) decan-1-ol-3-one	0.54	2-Propenoic acid, 3(4-methoxyphenyl), ethyl ester	0.26

33.	Crypton	2.47
34.	Cubenol	0.13
35.	Cuminal	0.23
36.	Cuminol	0.21
37.	Curcumene	0.16
38.	Cyclohexanone, 3-ethenyl	0.22
39.	Cyclohexene, 5-methyl-3-(1-methylethenyl)	0.10
40.	Damascone	0.32
41.	epi-13-Manool	0.11
42.	Eremophilene	0.10
43.	Isogeraniol	0.24
44.	Isolimonene	0.66
45.	Juniper camphor	0.43
46.	γ -Caryophyllene	0.53
47.	γ -Methylfuran	0.10
48.	Longipinocarvone	0.18
49.	γ -Pinene	6.31
50.	Methyl p-methoxycinnamate	5.02
51.	Methylvanillin	0.64
52.	Myrtanal	0.14
53.	Ocimene	0.83
54.	Pentadecyne	0.53
55.	Phellandral	0.42
56.	Pinocarvone	0.11
57.	Pseudo Limonene	0.15
58.	Sabinene	14.99
59.	tau-Muurolol	0.22
60.	Tetracyclo [6.3.2.0(2.5),0(1,8)] tridecan-9-ol, 4,4-dimethyl	0.29
61.	trans-Nerolidol	0.31
62.	trans-Pinocarveol	0.25
63.	Triquinacene, 1,4, bis(methoxy)	3.79
64.	Triquinacene, 1,4,7-tris (methoxy)	0.79

and its components identified. The high concentration of sabinene and triquinacene, 1,4-bis (methoxy) in leaf and rhizome oil makes it respectively potentially useful in the medicines

because they exhibit antibacterial activities. However, further study has to be conducted.

References

- Bin JI, Mohd YMS, Chin CB, Sim NL. Antifungal activity of the essential oils of nine Zingiberaceae species. *Pharm Biol.* 2003; 41: 392-97. <http://dx.doi.org/10.1076/phbi.41.5.392.15941>
- Giwanon R, Thubthimthed S, Rerk-am U, Sunthorntanasart T. Antimicrobial activity of terpinen-4-ol and sabinene. *Thai J Pharm Sci.* 2000; 24(Suppl): 27.
- Habsah M, Amran M, Mackeen MM, Lajis NH, Kikuzaki H, Nakatani H, Rahman A, Ghafar A, Ali AM. Screening of Zingiberaceae extracts for antimicrobial and antioxidant activities. *J Ethnopharmacol.* 2000; 72: 403-10. [http://dx.doi.org/10.1016/S0378-8741\(00\)00223-3](http://dx.doi.org/10.1016/S0378-8741(00)00223-3)
- Jeenapongsa R, Yoovathaworn K, Sriwatanakul KM, Pongprayoon U, Sriwatanakul K. Anti-inflammatory activity of (E)-1-(3,4-dimethoxyphenyl) butadiene from *Zingiber cassumunar* Roxb. *J Ethnopharmacol* 2003; 87: 143-48. [http://dx.doi.org/10.1016/S0378-8741\(03\)00098-9](http://dx.doi.org/10.1016/S0378-8741(03)00098-9)
- Jitoe A, Masuda T, Mabry TJ. Novel antioxidants, cassumunarin A, B, and C from *Zingiber Cassumunar*. *Tetrahedron.* 1994; 35: 981-84. [http://dx.doi.org/10.1016/S0040-4039\(00\)79944-8](http://dx.doi.org/10.1016/S0040-4039(00)79944-8)
- Kanjanapothi D, Soparat P, Panthong A, Tuntiwachwuttikul P, Reutrakul V. A uterine relaxant compound from *Zingiber cassumunar*. *Planta Med.* 1987; 53: 329-32. <http://dx.doi.org/10.1055/s-2006-962729>
- Kishore N, Dwivedi RS. Zerumbone: A potential fungitoxic agent isolated from *Zingiber cassumunar* Roxb. *Mycopathologia* 1992; 120: 155-59. <http://dx.doi.org/10.1007/BF00436393>
- Kuroyanagi M, Fukushima S, Yoshihira K, Natori S, Dechatiwongse T, Mihashi K, Nishi M, Hara S. Further characterization of the constituents of a Thai medicinal plant, *Zingiber cassumunar* Roxb. *Chem Pharm Bull.* 1980; 28: 2948-59.
- Masuda T, Jitoe A. Antioxidative and antiinflammatory compounds from tropical gingers: Isolation, structure determination, and activities of cassumunins A, B, and C, new complex curcuminoids from *Zingiber cassumunar*. *J Agricultural Food Chem.* 1994; 42: 1850-56. <http://dx.doi.org/10.1021/jf00045a004>
- Oliveros MB. Preformulation studies on terpinen-4-ol from *Zingiber purpureum* Rosc. In: Proceedings of the 2nd Symposium on the Family Zingiberaceae. Wu TL, Wu QG, Chen ZY (eds). Guangzhou, Zhongshan University Press, 1996, pp 180-86.
- Ozaki Y, Kawahara N, Harada M. Anti-inflammatory effect of *Zingiber cassumunar* Roxb. and its active principles. *Chem Pharm Bull.* 1991; 39: 2353-56.
- Panthong A, Kanjanapothi D, Niwatananant W, Tuntiwachwuttikul P, Reutrakul V. Anti-inflammatory activity of compound D ((E)-4-(3',4'-dimethoxyphenyl) but-3-en-2-ol) isolated from *Zingiber cassumunar*. *Phytomedicine* 1997; 4: 207-12.
- Panthong A, Kanjanapothi D, Niwatananun V, Tuntiwachwuttikul P, Reutrakul V. Anti-inflammatory activity of compounds isolated from *Zingiber cassumunar*. *Planta Med.* 1990; 56: 655. <http://dx.doi.org/10.1055/s-2006-961309>
- Piromrat K, Tuchinda M, Geadsomnuig S, Koysooko R. Antihistaminic effect of "Plai" (*Zingiber cassumunar* Roxb.) on histamine skin test in asthmatic children. *Siriraj Hospital Gazette* 1986; 38: 251-56.
- Pongprayoon U, Tuchinda P, Claeson P, Sematong T. Topical antiinflammatory activity of the major lipophilic constituents of the rhizome of *Zingiber cassumunar*. Part 1. The essential oil. *Phytomedicine* 1997; 3: 319-22.
- Pongprayoon U, Tuchinda P, Claeson P, Sematong T, Reutrakul V. Topical antiinflammatory activity of lipophilic constituents of the rhizome of *Zingiber cassumunar*. Part 2. Hexane extractives. *Phytomedicine* 1996; 3: 323.
- Tripathi P, Dubey NK, Shukla AK. Use of some essential oils as post-harvest botanical fungicides in the management of grey mould of grapes caused by *Botrytis cinerea*. *World J Microb Biotech.* 2008; 24: 39-46. <http://dx.doi.org/10.1007/s11274-007-9435-2>
- Taroeno JJ, Brophy JH, Zwaving. Analysis of the essential oil of *Zingiber cassumunar* Roxb. from Indonesia. *Fla Fra J* 1991; 6: 161-64. <http://dx.doi.org/10.1002/ffj.2730060214>
- Tuntiwachwuttikul P, Limchawfar B, Reutrakul VPO, Kusamran K, Byrne LT. Syntheses of some constituents of *Zingiber cassumunar*. *Aust J Chem.* 1980; 33: 913-16.
- Tuntiwachwuttikul P, Pancharoen O, Jaipetch T, Reutrakul V. Phenylbutanoids from *Zingiber cassumunar*. *Phytochemistry* 1981; 20: 1164-65. [http://dx.doi.org/10.1016/0031-9422\(81\)83058-0](http://dx.doi.org/10.1016/0031-9422(81)83058-0)
- Wasuwat S, Wanisorn P, Mahintorntep B, Kuwaono K, Sanghirun S. Studies on antimicrobial and antifungal activities of terpinen-4-ol extracted from *Zingiber cassumunar* Roxb. Thailand Institute of Scientific and Technological Research 1989, Research Project No. 30-32/Rep. No.1 (In Thai).

