

BJP

Bangladesh Journal of Pharmacology

Research Article

**Detect the LD50 of a Preservative
Sodium Nitrite (E250) in Female Rats**

Detect the LD50 of a Preservative Sodium Nitrite (E250) in Female Rats

Asst. Lec. Zainab Majid Mohammed¹, Asst. Lec. Zainab Abdul Abbas Nasser², Asst. Lec. Rana Hasan Hadi Albuissa³, Asst. Lec. Maryam Isam Abdullah Abdalzahrh Alnaser⁴

¹Biological, Al Kufa University, Work in Al-Kafeel University, zainab.majid@alkafeel.edu.iq

²Biological, Al Kufa University, zainaba.

³Biology, University of Alkufa, Babylon Education Directorate, ranah.

⁴Molecular Biology, Department of Analytical Science, University of Alkufa,

Article Info

Cite this article:

Asst. Lec. Zainab Majid Mohammed, Asst. Lec. Zainab Abdul Abbas Nasser, Asst. Lec. Rana Hasan Hadi Albuissa, Asst. Lec. Maryam Isam Abdullah Abdalzahrh Alnaser. Detect the LD50 of a Preservative Sodium Nitrite (E250) in Female Rats. Bangladesh J Pharmacol. 2026; 21: 1-7.

Key words Sodium nitrite, E250, median lethal dose (LD50), Rattus rattus, toxicology, preservative.

Abstract

The present study was conducted to detect the LD50 (median lethal dose) and toxicological effects of a preservative, sodium nitrite (E250), oral administration in female rats. We performed this experiment at the Animal House Facility, College of Education for Girls, University of Kufa from October 11 to December 16, 2022. Adult female albino rats (*Rattus rattus*), around 13 weeks of age and weighing between 200 g, were used. On the other hand, the study has precisely estimated LD50 and toxic dose of sodium nitrite. Twenty-eight female rats were divided into seven groups, each four animals. The variant doses of sodium nitrite groups were administered orally at 20, 30, 45, 70, 100, 130 and 160 mg/kg body weight using a gastric gavage tube once a day. The differences in the acute oral toxicity of 70, 100, 130 and 160 mg/kg body weight were significant for female rats dying within last period after dosing between 24–48 hours (the lethal concentrations). A 45 mg/kg body weight dose was the LD50 recorded in female rats, with mortality of half the rats within 48–72 hours. Nevertheless, this dosage eventually killed those remaining rats after a minimum of seven days. Nonetheless, both 30 and 20 mg/kg body weight doses were non-lethal and non-toxic. Therefore, the use of doses under 20 mg/kg (10 and 15 mg/kg body weight) was considered safe for preservative in products. The study its for the disease of sodium nitrite (E250) in *Rattus rattus* female LD50 45mg/Kg..

Introduction

Food additives, including chemical preservatives, can pose health risks when used in excessive quantities. Consequently, the United States Food and Drug Administration (FDA) has established stringent regulations for the approval of various synthetic food additives (Mister and Hathcock, 2012). Additionally, the World Health Organization (WHO) and the European Food Safety Authority (EFSA) have collaborated to develop a program for the evaluation and designation of food additives.

Aims of the Study

The present study aims to determine the median lethal dose (LD50) and assess the toxicity of the preservative sodium nitrite (NaNO_2 , E250) in female

albino rats (*Rattus rattus*)

Literature Review

Preservatives in Food Systems

Preservatives are organic or inorganic compounds added to foods in order to prevent microbial growth and undesirable chemical changes that can extend the shelf life of a food product (Sebranek and Bacus, 2007). One of the most commonly used preservatives is sodium nitrite (NaNO_2 ; E250), which has been predominantly used to prevent lipid oxidation in meat preservation (Sindelar & Milkowski, 2012). Sodium nitrite (E250): The Joint Expert Committee on Food Additives (JECFA, 2002) again re-evaluated sodium nitrite and set an ADI



from 0.06 to 0.07 mg/kg body weight per day for almost all population groups. Methemoglobinemia, a condition in which nitrite binds with hemoglobin to form methemoglobin, and blocks the binding site of oxygen on it, thus preventing oxygen uptake by blood is one of the most important toxicological effects of sodium nitrite in the blood. Symptoms include cyanosis of the lips and peripheries, as well as diarrhea which is common in infants, while methemoglobin levels > 70% in blood can be fatal due to anoxia. This syndrome is also well recognized, known as "blue baby" (frequently), (Sanchez et al., 2001). The brain, which uses more oxygen than any other organ in the body, is especially sensitive to lack of oxygen and can begin to die within five minutes after oxygen supply has been cut. This then can cause problems such as lack of attention, stability imbalance, and lack of mobility function (Kaur et al., 2005). In addition, nitrite used for preservation or colorant has been reported to be linked with occurrence of cancer, such as gastric and esophageal cancer, colorectal cancer, liver cancer, bladder TCCs and thyroid carcinoma (Song et al. 2015). For example, long-term intake of sodium nitrite has been shown to cause adverse effects on the male reproductive system via a consequential decrease in testosterone and follicle-stimulating hormone (FSH) levels. Such changes lead to lower numbers of viable spermatozoa, higher proportions of dead and morphologically abnormal sperm, subsequently decreasing the fertility potential in males (Farias et al., 2005). However, sodium nitrite is capable of causing renal dysfunction by increasing levels of blood urea and creatinine, suppression in proximal tubule reabsorption and also triggering glomerular hyperfiltration (Helal et al., 2000). The studies demonstrated decreased levels of antioxidants caused by sodium nitrite exposure, inhibiting mechanisms that lead to suppression of steroid hormone production in the ovaries and adrenal glands, with lower enzymatic antioxidant activity on pituitary tissues as well as on ovarian hormone synthesis. Unfortunately, this results in ovarian follicle loss and impaired female fertility (Creanga et al., 2008). Studies by EL-Nabarawy et al. Kiani et al. (2017) revealed that the treatment with 75 mg/kg sodium nitrite in males resulted in an increase of free radical formation, lipid peroxidation and creatinine as well as urea levels which may lead to kidney and liver tissue damage, subsequently can cause hepatic dysfunction(2020).

Additives Food

Food additives are chemical substances of natural or synthetic origin, which are added to foodstuffs specifically for their technical and/or organoleptic (Sensory) effect in the finished product (Linkie et al. According to the report of (Chazelas et al.), food additives are categorized according to their function in food. (2020), into the following categories:

Coloring agents

Acidity regulators and anti-caking agents

Flavoring agents

Thickeners, stabilizers, and emulsifiers

Firming agents and foaming agents

Preservatives

What are Preservatives—Preservative is a chemical component that is added to food products for the prevention of spoilage from microbial growth, enzymatic activity or oxidation. They are important in the food, pharmaceutical and cosmetic industries through prevention of bacterial growth and bio-enrichment (Al-Dhafer, 2008).


Types of Preservatives

Natural Preservatives: For ex-salt, spices, sugar, oils and etc. used widely in the production of jams, juices and pickles (Smith and Hong 2011).

Synthetic Preservative: These are used to reduce the chances of chemical and biological spoilage of food, although there is an amount that needs to be monitored as these preservatives are known to cause cancer if they exceed a safe level. Based on both chemical composition and mechanism of action they can further be classified as:

Organic Preservatives: These are sorbic acid, benzoic acid and acetic acid.

Inorganic Preservatives: like nitrates (NO_3^-) and nitrites which is (NO_2^-). International designator E250 Chemical structure of sodium nitrite Sodium nitrite is one type of inorganic preservative. Daily dosing of sodium nitrite is safe in humans and accepted internationally at a dose approximate to 10 mg/kg body weight (Carocho et al., 2014).



Chemical Composition of Sodium Nitrite (Carocho,etal.,2014)

Table 1: Chemical and Physical Properties of Sodium Nitrite (NaNO_2)

Property	Description
Chemical Formula	NaNO_2

Molecular Weight	68.9953 g/mol
Melting Point	271 °C
Density	2.17 g/cm ³
Boiling Point	115 °C (decomposes)
Classification	Inorganic compound, corrosion inhibitor
Solubility	Soluble in water and ethanol
IUPAC Name	Sodium nitrite
Appearance	White to yellowish crystals
Alternative Names	E250
Solubility in Water	82 g/100 mL water at 20 °C
Taste	Slightly salty
Decomposition Temperature	320 °C (608 °F)

Sources of Sodium Nitrite

Drinking Water: About 25% of salivary nitrate is metabolized to nitrite by oral microflora, especially the bacteria located in the dorsal surface of the tongue. Gilchrist et al. (2010) found that the intestine absorbs this nitrite to be used later. Nitrate to nitrite reduction in the oral cavity is probably the major source of human exposure to nitrite especially from dietary sources (EFSA, 2008).

Plants and Soil: Source of nitrates — green leafy vegetables, root vegetables. The use of fertilizers leads to increase in Nitrogen and the consumption of organic material produced a result a form changing it into N which is used by plants to form nitrate, and this result for many nitration of vegetable (WHO 2008).

Antioxidants

Sodium Nitrite Source and Application in Regulation level Sodium nitrite at 20mg/kg decreased the lipid oxidation of meat product (Al-Shuibi et al., 2002). A study reported reducing the concentration down to 50 mg/kg and still achieving effectiveness as an antioxidant in beef and poultry (Sebranek, J., Bacus, J.N., 2007) Sodium nitrite in combination with salts is added to processed meats as a safety enhancer since it inhibits the growth of pathogenic microorganisms (Mulvey et al. 2010). Concentrations of nitrite up to 45–119 mg/kg have been shown to stabilize color in meats (Sebranek et al., 2007) with an ideal concentration for maximal color stability between 10 and 20 mg/kg (Ranken, 2000).

Sodium nitrite is also employed in the commercial food sector as a colour fixative in confectionery, dairy goods, jams, preserved meats, gelatin, dry soft drink mixes and canned foods at recommended levels ranging from 5–20 mg/kg to give a red-bound coloration to meats (Pradhan et.al 2019). This compound is used in food and beyond food applications (pharmaceutical, cosmetic) including aqueous pharmaceutical solution, tablets (0.1 g or 110 mg sodium nitrite), capsules (110 or 30mg NaNO₂), toothpaste, cosmetics as well as industrial uses — ink production [5]. Despite the forgoing effects, CNV is also a vasodilator; when used in human and veterinary medicine it has been utilized to treat various vascular conditions as well (Liu et al., 2020). For this purpose, intramuscular injections of sodium nitrite (1–10 mg/kg) have been proven to be effective at mitigating the adverse effects related to exposure to chlorine gas as well as in treating cyanide and carbon monoxide poisoning, especially when dealing with life-threatening circumstances (Lutfiya Miller et al., 2021). International Regulations for Sodium Nitrite

The percentage of nitrite and nitrate must meet a given order of magnitude in processed meats, which are used to stabilize the color and prevent the growth of bacteria. The Joint Expert Committee on Food Additives (JECFA) in the Food and Agriculture Organization of the United Nations/World Health Organization (WHO)[and8] have established an acceptable daily intake as 0.06–0.07 mg/kg body weight for a 60-kilogram- (130 lb-) adult, equal to 4.2 mg/day. The maximum allowable level of sodium nitrite that can be added to processed meats and fish is 200 mg/kg in the United States (FDA [Food and Drug Administration], 2017.)

Materials and Methods

Animals Labrotary

Albino female adult rats (*Rattus rattus albinus* Sprague-Dawley) aged between 10 and 13 weeks, weighing between 200-220 g were used in the experiment. Rats were obtained from the Animal House Facility of the College of Sciences at the University of Kufa, and their non-pregnancy, disease-free status and that they had not been previously used in any experimental designs was further confirmed. Animals were provided with normal feed and drinking water through bottles. The animals were kept for 1 week in the laboratory under standard conditions before the initiation of experiments. Experimental rats: The female rats were housed in special dosing cages and the rat sprayers wrote down the start date of the experiment and their name. The cages were provided water and feed, and dosing was carried out in a rat proof oral gavage tool shown in the figure..



Gavage Dosing Procedure

Design of experiments

The intention of this experiment is designed to estimate the median lethal dose (LD50) of sodium nitrite (NaNO_2) in female rats at day 1. The study was conducted on 28 adult female albino rats (*Rattus rattus* Sprague-Dawley strain). Animals and treatment Seven female Wistar rats were divided into seven group with four animal each. Rats received a 0.1 mL sodium nitrite through a metal stomach tube, orally once daily (Sarkar et al., 2005) as follow

Group 1: [4] female rats dosed by gavage with sodium nitrite at a concentration of 160 mg/kg body weight.

Groups 2: Four female rats received via oral intubation with Sodium nitrite at a concentration of 130 mg/kg B W,

Group 3 (four female rats were injected with 100 mg/kg body weight of sodium nitrite.)

Group 4: Four female rats were administered the preservative sodium nitrite at a concentration of 70 mg/kg body weight.

Group 5: Four female rats were administered a preservative of sodium nitrite(45 mg/kg body weight.)

Group 6: Four normal female rats administered the preservative sodium nitrite at dose 30 mg/kg body weight.

Group 7. 4 female rats — preservative = sodium nitrite at a dose level of 20 mg/kg body weight.

Results

Results: All female rats injected with sodium nitrite at doses of 160, 130, 100 and 70 mg/kg body weight died within the first day of treatment. Those levels were classed as toxic, and mortal to female rats. On the other hand, half of the rats treated with 45 mg/kg body weight died within 48–72 hours and all others in the next 7–10 days. Therefore, this dose (45 mg/kg) was the LD50 of sodium nitrite for female rats. In light of this finding, lower doses (30 and 20 mg/kg body weight) were also

applied and the results showed that these concentrations are not lethal to female rats, since all treated animals survived. Consequently, further female rat dosing studies were carried out using doses below the LD50 and specifically lower than 20 mg/kg body weight.

Table 2: Determination of the Median Lethal Dose (LD50) and Toxicity of Sodium Nitrite in Female Rats

Concentration (Dose)	Treated Rats	Number of Rats Died Within 24 Hours Post-Dosing	Number of Rats Died After 24 Hours Post-Dosing	Effects on Female Rats
160 mg/kg body weight	4	4	0	Let hal
130 mg/kg body weight	4	4	0	Let hal
100 mg/kg body weight	4	4	0	Let hal
70 mg/kg body weight	4	3	1	Let hal
45 mg/kg body weight	4	2	2	Let hal
30 mg/kg body weight	4	0	1	To xic
20 mg/kg	4	0	0	To xic

body weight				
-------------	--	--	--	--

Discussions

In this experiment, all female rats treated with sodium nitrite concentrations of 160, 130, 100 or 70 mg/kg body weight on the 1st day of died within a day after exposure. The toxic or lethal doses (concentration) for female rats were therefore recorded. Female rats killed with 45 mg/kg body weight showed in half of the animals death after 48-72 hours, all others died within 7–10 days. As a result, 45 mg/kg was regarded as the median lethal dose (LD50) of sodium nitrite for female rats. Considering the above result, lower doses of 30 and 20 mg/kg body weight were applied and such concentrations were not lethal as all female rats treated with an equivalent dosage survived. These observed effects could be associated with sodium nitrite interacting with the nitric oxide (NO) present in the organism, causing vasodilatation and vascular smooth muscle relaxation, resulting in a fall off of oxygen delivery leading to severe tissue injury at organs such as kidneys and liver, respectively followed by death (Arnold et al., 2020). Other research also links sodium nitrite to cerebral hypoxia and brain tissue damage thanks to the way it prevents your bloodstream from getting oxygen. Mild hypoxia results in symptoms that include ataxia and short-term memory loss, whereas severe or prolonged hypoxia can lead to coma, seizures, and cerebral death (Cymerman 2001). This is consistent with the results of Al-Hasnawi (2021) who found that a single dose of 60 and 100 mg/kg body weight caused mortality in male and female albino rats, while repeated doses of 10 to 30 mg/kg body weight did not cause any death. So, too, did the present study demonstrate that doses of 20–30 mg/kg body weight (given to female rats) were not lethal..

References

- [1] Al-Shuib A., Al-Abdullah B. (2002). "Substitution of nitrite by sorbate and the effect on properties of mortadella". *Meat Science.*, 62: 473-478.
- [2] Arnold, J. T., Lloyd, A. B., Bailey, S. J., Fujimoto, T., Matsukate, R., Takayanagi, M., Nishiyasu, T., and Fujii, N. (2020). The nitric oxide dependence of cutaneous microvascular function to independent and combined hypoxic cold exposure. *Journal of Applied Physiology.*
- [3] Carocho, M; Barreiro, M. F; Morales, P; and Ferreira, I. C. (2014). Adding molecules to food, pros and cons: A review on synthetic and natural food additives. *Comprehensive Reviews in Food Science and Food Safety.*, 13(4): 377-399.
- [4] Chazelas, E; Deschasaux, M; Srour, B; Kesse-Guyot, E; Julia, C., Allès, B; Druetne-Pecollo, N; Galan, P; Hercberg, S; and Latino-Martel, P. (2020). Food additives: distribution and co-occurrence in 123,000 food products of the French market. *Scientific Reports*, 10(1): 1–15.
- [5] EFSA (European Food Safety Authority) .(2008). Nitrate in vegetables—Scientific Opinion of the Panel on Contaminants in the Food chain. *EFSA J*, 689:1–79
- [6] EFSA. (2011). Panel on food additives and nutrient sources added to food (ANS); Scientific opinion on the reevaluation of butylated hydroxyanisole–BHA (E 320) as a food additive. *EFSA J* 9(10):2392 .
- [7] El-Nabarawy, N. A., Gouda, A. S., Khattab, M. A., and Rashed, L. A. (2020). Effects of nitrite graded doses on hepatotoxicity and nephrotoxicity, histopathological alterations, and activation of apoptosis in adult rats. *Environ. Sci. Poll. Res:1–14.Explants. Methods. Mol. Biol.* ..2092:65-75.
- [8] Farias, J. G., Bustos-Obregón, E., Orellana, R., Bucarey, J. L., Quiroz, E., and Reyes, J. G. (2005). Effects of chronic hypobaric hypoxia on testis histology and round spermatid oxidative metabolism. *Andrologia.*, 37(1): 47–52.
- [9] Helal, E.G.E; Zahkoui, S.A, M and Mekawy, H.A. (2000): Effect of some food colours (synthetic and natural products) on liver and kidney functions of young albino rats. *Egyptian J. of Hospital. Med.*
- [10] JECFA Joint FAO/WHO Expert Committee on Food Additives, (2002). Evaluation of certain food additives and contaminants. Fifty-ninth report of JECFA. Joint FAO/WHO Expert Committee on Food Additives, WHO Technical Reports Series No. 913: 20–32.
- [11] Kaur, C; Sivakumar .V;Singh. G; Singh .J; Ling EA (2005).Response of Purkinje neurons to hypobaric hypoxic exposure as shown by alteration in expression of glutamate receptors, nitric oxide synthases and calcium binding proteins. *Neuroscience.*,135(4):1217-1229.
- [12] Lide, D. (2005). *Handbook of Chemistry and Physics*, 86th edn. CRC. Taylor and Francis.
- [13] Linke ,BGO. (2018).Casagrande TAC and Cardoso LAC., *African. Biotech.*,17(10):306-310
- [14] Pradhan, H; Poudel, S; Bokkhim, H; Rai, K. P. and Karn, S. K. (2019) Status of Sodium Nitrite in Meat and Meat Products Available in the Market of Kathmandu, Nepal. *J. Food Sci. Technol Nepal*, 11, 47–50.
- [15] Ranken M. (2000). "Handbook of meat product technology (pp. 54-62). Blackwell Science 12.
- [16] Sebranek JG,and Bacus JN. (2007). Meat products without direct addition of nitrate or nitrite: what are the issues? *Meat Sci* 77:136–47.

- [17] Sindelar JJ, and Milkowski AL.(2012). Human safety controversies surrounding nitrate and nitrite in the diet. Nitric Oxide 26:259–66. http://www.who.int/water_sanitation_health/dwq/gdwq0506.pdf.
- [18] Song, P., Wu, L. and Guan, W. Dietary(2015). Nitrates, Nitrites, and Nitrosamines Intake and the Risk of Gastric Cancer: A Meta-Analysis. Nutrients 7, 9872–9895
- [19] WHO, (World Health Organization) (2008). "Guidelines for Drinking-waterQuality
- [20] Al-Hasnawi, Maytham Ali Karim. (2021). The aqueous extract of garden cress (*Lepidium sativum*) seeds on some histological and functional parameters against the toxic effects induced by sodium nitrite in male albino rats, Master's thesis, College of Education for Pure Sciences, University of Karbala..
-